

In certain embodiments, the culture is contacted with a steroid or corticosteroid such as, for example, hydrocortisone, deoxyhydrocortisone, fludrocortisone, prednisolone, methylprednisolone, prednisone, triamcinolone, dexamethasone, betamethasone and paramethasone. See, generally, The Merck Manual of Diagnosis and Therapy, 15th Ed., pp. 1239-1267 and 2497-2506, Berkow et al., eds., ~~Rahay~~ Rahway, N.J., 1987).

In the claims:

For the convenience of the Examiner, all claims being examined, whether or not amended, are presented below.

C4 55. **(Amended)** A method for preparing a substantially pure non-adherent population of progenitor cells comprising:
obtaining a cell suspension from an animal tissue selected from pancreatic tissue or pancreatic ductal tissue, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about 75% pure.

C5 57. **(Amended)** A method for preparing a substantially pure non-adherent population of progenitor cells comprising:
obtaining a cell suspension from an animal tissue selected from pancreatic tissue or pancreatic ductal tissue, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about 80% pure.

58. **(Twice Amended)** The method of claim 79, wherein said non-adherent population of progenitor cells is at least about 75% pure.

59. **(Twice Amended)** The method of claim 79, wherein said non-adherent population of progenitor cells is at least about 80% pure.

60. **(Reiterated)** The method of claim 57, wherein said non-adherent population of progenitor cells is at least about 90% pure.

61. **(Reiterated)** The method of claim 57, wherein said animal tissue is obtained from a mammalian organ.

63. **(Reiterated)** The method of claim 55, wherein said cell suspension is obtained by mechanical disruption of said animal tissue.

64. **(Reiterated)** The method of claim 55, wherein said cell suspension is obtained by enzymatic disruption of said animal tissue.

65. **(Reiterated)** The method of claim 55, wherein said growth factor preparation comprises at least one of epidermal growth factor, transforming growth factor, hepatocyte growth factor, fibroblast growth factor, leukemia inhibitory factor, insulin-like growth factor, and platelet-derived growth factor.

66. **(Reiterated)** The method of claim 57, wherein said substantially pure non-adherent progenitor cells are floating cells.

68. **(Thrice Amended)** The method of claim 79, wherein said substantially pure non-adherent progenitor cells form a homotypic cell sphere.

69. **(Amended)** A method for preparing a substantially pure non-adherent population of progenitor cells comprising:
providing an animal tissue selected from pancreatic tissue or pancreatic ductal tissue;

disrupting said animal tissue so as to obtain a cell suspension comprising at least one progenitor cell; and

allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,

thereby obtaining a substantially pure non-adherent progenitor cell population at least about 75% pure.

C6
C7
71. (Amended) The method of claim 55, 69 or 79, wherein said non-adherent progenitor cell population expresses Nestin.

72. (Amended) The method of claim 55, 69 or 79, wherein said non-adherent progenitor cell population expresses at least one of c-kit and Sca.

73. (Reiterated) The method of claim 55 or 69, wherein said non-adherent progenitor cell population under proper conditions can give rise to cells that express a marker selected from Pdx-1, glucagon, and insulin.

C8
74. (Amended) A composition comprising the substantially pure nonadherent progenitor cell population obtained by the method of claim 55, 69, 79 or 81.

75. (Reiterated) The composition of claim 74, wherein the substantially pure nonadherent progenitor cell population expresses a marker selected from Nestin, c-kit, and Sca.

76. (Reiterated) The composition of claim 74, wherein the substantially pure nonadherent progenitor cell population under proper conditions can give rise to cells that express a marker selected from Pdx-1, glucagon, and insulin.

C9
77. (Amended) The method of claim 79 or 81, wherein said substantially pure non-adherent population of progenitor cells is at least about one thousand-fold enriched from said animal cell suspension.

78. (Amended) The method of claim 79, wherein said substantially pure non-adherent population of progenitor cells is at least about one hundred-fold enriched from said animal cell suspension.

Please add the following new claims:

79. (NEW) A method for preparing a substantially pure non-adherent population of progenitor cells comprising:

obtaining a cell suspension from an animal tissue selected from liver tissue or dermis, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about 50% pure.

80. (NEW) A method for preparing a substantially pure non-adherent population of progenitor cells comprising:

obtaining a cell suspension from an animal tissue selected from liver tissue or dermis, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about 60% pure.

81. (NEW) A method for preparing a substantially pure non-adherent population of progenitor cells comprising:

providing an animal tissue selected from liver tissue or dermis;
disrupting said animal tissue so as to obtain a cell suspension comprising at least one progenitor cell; and

allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,

thereby obtaining a substantially pure non-adherent progenitor cell population at least about one hundred-fold enriched from said animal cell suspension.

The claims presented above incorporate changes as indicated by the marked-up versions below.

55. (Amended) A method for preparing a substantially pure non-adherent population of progenitor cells comprising:
obtaining a cell suspension from an animal tissue selected from pancreatic tissue; or pancreatic ductal tissue, ~~liver tissue, and dermis~~, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about ~~50~~ 75% pure.

57. (Amended) A method for preparing a substantially pure non-adherent population of progenitor cells comprising:
obtaining a cell suspension from an animal tissue selected from pancreatic tissue; or pancreatic ductal tissue, ~~liver tissue, and dermis~~, wherein said cell suspension comprises at least one progenitor cell;
treating the cell suspension with a growth factor preparation; and
allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,
thereby obtaining a substantially pure non-adherent progenitor cell population that is at least about ~~60~~ 80% pure.

58. **(Twice Amended)** The method of claim 79 ~~57~~, wherein said non-adherent population of progenitor cells is at least about ~~70~~ 75% pure.

59. **(Twice Amended)** The method of claim 79 ~~57~~, wherein said non-adherent population of progenitor cells is at least about 80% pure.

68. **(Thrice Amended)** The method of claim 79 ~~57~~, wherein said substantially pure non-adherent progenitor cells form a homotypic cell sphere.

69. **(Amended)** A method for preparing a substantially pure non-adherent population of progenitor cells comprising:

providing an animal tissue selected from pancreatic tissue; or pancreatic ductal tissue; ~~liver tissue, and dermis;~~

disrupting said animal tissue so as to obtain a cell suspension comprising at least one progenitor cell; and

allowing proliferation of said at least one progenitor cell such that a substantially pure non-adherent progenitor cell population is obtained,

thereby obtaining a substantially pure non-adherent progenitor cell population at least about 75% ~~pure one hundred fold enriched from said animal cell suspension.~~

71. **(Amended)** The method of claim 55, ~~or~~ 69 or 79, wherein said non-adherent progenitor cell population expresses Nestin.

72. **(Amended)** The method of claim 55, ~~or~~ 69 or 79, wherein said non-adherent progenitor cell population expresses at least one of c-kit and Sca.

74. **(Amended)** A composition comprising the substantially pure nonadherent progenitor cell population obtained by the method of claim 55, ~~or~~ 69, 79 or 81.